



Acetaminophen (Tylenol) Dosing Chart

Date: _____ Child's Weight: _____

Give every 4-6 hours as needed, no more than 5 times in 24 hours.

*** The old infant drops (80mg/0.8ml) are no longer available. The new infant drops have the same concentration as the children's liquid *****

Weight	Milligram Dosage	Infant Drops AND Children's Liquid 160mg/5ml 5 ml = 1 teaspoon	Chewable Tablets 80mg	Jr. strength caps or chews 160mg	Adult Tablets 325mg
5-8 lbs	40 mg	1/4 teaspoon (1.25 ml)			
9-10 lbs	60 mg	1/3 teaspoon (1.8 ml)			
11-16 lbs	80 mg	1/2 teaspoon (2.5 ml)			
17-21 lbs	120 mg	3/4 teaspoon (3.75 ml)			
22-26 lbs	160 mg	1 teaspoon (5ml)	2 tablets	1 tablet	
27-32 lbs	200 mg	1 1/4 teaspoon (6.25 ml)	2 1/2 tablets	1 tablet	
33-37 lbs	240 mg	1 1/2 teaspoon (7.5 ml)	3 tablets	1 1/2 tabs	
38-42 lbs	280 mg	1 3/4 teaspoon (8.75 ml)	3 1/2 tablets	1 1/2 tabs	
43-53 lbs	320 mg	2 teaspoon (10 ml)	4 tablets	2 tablets	1 tablet
54-64 lbs	400 mg	2 1/2 teaspoon (12.5 ml)	5 tablets	2 1/2 tablets	1 tablet
65-75 lbs	480 mg	3 teaspoon (15 ml)	6 tablets	3 tablets	1 1/2 tablets
76-86 lbs	560 mg	3 1/2 teaspoon (17.5 ml)	7 tablets	3 1/2 tablets	1 1/2 tablets
87-95 lbs	640 mg	4 teaspoon (20 ml)	8 tablets	4 tablets	2 tablets
>95 lbs		GIVE	ADULT	DOSE	

Important Note:

1. Always use a proper measuring device. For example:
 - a. When giving infant drops, use only the dosing device (dropper or syringe) enclosed in the package.
 - b. When giving children's suspension or liquid, use the dosage cup enclosed in the package. (kitchen spoons are not accurate measures)

West End Location
4687 Pouncey Tract Road
Glen Allen, VA 23059
Phone: 804-422-KIDS (5437)
Fax: 804-474-9071

Southside Location
5021 Craig Rath Blvd, Bldg. IV
Midlothian, VA 23112
Phone: 804-592-KIDS (5437)
Fax: 804-592-2406

Mechanicsville Location
8356 Bell Creek Road
Mechanicsville, VA 23116
Phone: 804-559-KIDS (5437)
Fax: 804-277-8140

Stafford Location
20 Doc Stone Road
Stafford, VA 22556
Phone: 540-602-7766
Fax: 540-699-2383





Ibuprofen (Motrin, Advil) Dosing Chart

Date: _____ Child's weight: _____

Give every 6-8 hours, as needed, no more than 4 times in 24 hours

**** Not for use in children under 6 months of age ****

Ibuprofen infant drops: 1/3 syringe = 0.625 ml, 2/3 syringe = 1.25 ml, 1 syringe = 1.875 ml

West End Location
4687 Pouncey Tract Road
Glen Allen, VA 23059
Phone: 804-422-KIDS (5437)
Fax: 804-474-9071

Southside Location
5021 Craig Rath Blvd, Bldg. IV
Midlothian, VA 23112
Phone: 804-592-KIDS (5437)
Fax: 804-592-2406

Mechanicsville Location
8356 Bell Creek Road
Mechanicsville, VA 23116
Phone: 804-559-KIDS (5437)
Fax: 804-277-8140

Stafford Location
20 Doc Stone Road
Stafford, VA 22556
Phone: 540-602-7766
Fax: 540-699-2382

Weight	Milligram Dosage	Infant Drops 50mg/1.25ml	Children's Liquid 100mg/5ml	Chewable Tablets 50mg	Junior strength 100mg	Adult Tablets 200mg
9-10 lbs >6 mo.	25 mg	1/3 syringe (0.625 ml)				
11-16 lbs	50 mg	2/3 syringe (1.25 ml)	1/2 teaspoon (2.5ml)			
17-21 lbs	75 mg	1 syringe (1.875 ml)	3/4 teaspoon (3.75ml)			
22-26 lbs	100 mg	1 1/3 syringe (1.875ml + 0.625ml)	1 teaspoon (5 ml)	2 tablets	1 tablet	
27-32 lbs	125 mg	1 2/3 syringe (1.875ml + 1.25ml)	1 1/4 teaspoon (6.25 ml)	2 1/2 tablets	1 tablet	
33-37 lbs	150 mg	2 syringes (2 x 1.875ml)	1 1/2 teaspoon (7.5 ml)	3 tablets	1 1/2 tablets	
38-42 lbs	175 mg	2 1/3 syringes (2 x 1.875ml + 0.625 ml)	1 3/4 teaspoon (8.75 ml)	3 1/2 tablets	1 1/2 tablets	
43-53 lbs	200 mg	2 2/3 syringes (2 x 1.875 ml + 1.25 ml)	2 teaspoon (10 ml)	4 tablets	2 tablets	1 tablet
54-64 lbs	250 mg	Use liquid or tablets.	2 1/2 teaspoon (12.5 ml)	5 tablets	2 1/2 tablets	1 tablet
65-75 lbs	300 mg		3 teaspoon (15 ml)	6 tablets	3 tablets	1 1/2 tablets
76-86 lbs	350 mg		3 1/2 teaspoon (17.5 ml)	7 tablets	3 1/2 tablets	1 1/2 tablets
87-95 lbs	400 mg		4 teaspoon (20 ml)	8 tablets	4 tablets	2 tablets
> 95 lbs		GIVE	ADULT	DOSE		

Important Note:

1. Always use a proper measuring device. For example:
 - a. When giving infant drops, use only the dosing device (dropper or syringe) enclosed in the package.
 - b. When giving children's suspension or liquid, use the dosage cup enclosed in the package. (kitchen spoons are not accurate measures)

